Package 'demofit'

November 7, 2026

Title Parametric Mortality Curve Fitting and Mortality Forecasting Tools

Version 0.1.6

Author Jackie Li [aut, cre, cph]

Maintainer Jackie Li <jackieli@smu.edu.sg>

Description Provides tools for fitting parametric mortality curves. Implements multiple optimisation strategies to enhance robustness and stability of parameter estimation. Offers tools for forecasting mortality rates guided by mortality curves.
For modelling details see:
Tabeau (2001) <doi:10.1007/0-306-47562-6_1>,
Renshaw and Haberman (2006) <doi:10.1016/j.insmatheco.2005.12.001>,
Cairns et al. (2009) <doi:10.1080/10920277.2009.10597538>,
Li and Lee (2005) <doi:10.1353/dem.2005.0021>.

License GPL-3

Encoding UTF-8

RoxygenNote 7.3.3

Imports forecast, minpack.lm, MortalityLaws, NlcOptim, stats, MASS, graphics, grDevices

Suggests knitr, rmarkdown, ggplot2, scales

VignetteBuilder knitr

NeedsCompilation no

Repository CRAN

Date/Publication 2026-11-07 21:50:02 UTC

Contents

APCS 2
CAES 4
CBDCS 5
CBDQCS 6
CBDS 8

CC . . . 9
CFM2S . . . 10
CFMS . . . 11
curveS3 . . . 13
ENI . . . 13
ENS . . . 15
FCS . . . 16
LCLS . . . 17
LCS . . . 19
MC . . . 20
modelS3 . . . 22
mortalitymatrix . . . 23
PAPCS . . . 24
PCAES . . . 25
PCBDCS . . . 27
PCBDQCS . . . 28
PCBDS . . . 29
PCFM2S . . . 31
PCFMS . . . 32
PFCS . . . 34
PLCS . . . 36
PRHS . . . 37
RHS . . . 39
SEN . . . 40
STARS . . . 42

Index **44**

`APCS` *Age-period-cohort model*

Description

Fits and forecasts mortality rates using age-period-cohort model.

Usage

```
APCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

`x`	vector of ages.
`M`	matrix of mortality rates (rows as years and columns as ages).
`curve`	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2,

	vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The age-period-cohort (APC) model is specified as

$ln(m_{x,t}) = \alpha_x + \kappa_t + \gamma_{t-x} + \epsilon_{x,t}$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c. Constraints include sum of κ_t is zero and and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class APCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Bray, I. (2002). Application of Markov chain Monte Carlo methods to projecting cancer incidence and mortality. Journal of the Royal Statistical Society Series C, 51(2), 151-164.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- rep(1/30,30)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- APCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CAES *Common age effect model*

Description

Fits and forecasts mortality rates of two populations using common age effect model.

Usage

```
CAES(x, M1, M2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M1	matrix of mortality rates of population 1 (rows as years and columns as ages).
M2	matrix of mortality rates of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common age effect (CAE) model is specified as

$ln(m_{x,t,i}) = \alpha_{x,i} + \beta_x \kappa_{t,i} + \epsilon_{x,t,i}$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{t,i}$. Constraints include sum of β_x is one and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class CAES with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Kleinow, T. (2015). A common age effect model for the mortality of multiple populations. Insurance: Mathematics and Economics, 63(C), 147-152.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
b <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
k1 <- c(8.68,8.34,7.99,6.87,8.18,5.73,4.83,5.20,2.74,3.22,
2.99,1.59,1.67,-0.65,-0.39,-1.07,-0.95,-2.78,-3.46,-2.45,
-4.12,-4.66,-4.98,-4.58,-6.30,-4.39,-5.56,-6.52,-8.26,-6.92)
k2 <- c(11.81,11.01,10.59,10.40,9.75,8.15,6.07,6.45,4.60,4.57,
4.15,1.49,1.77,-1.08,-1.44,-0.96,-1.66,-2.25,-4.67,-4.62,
-4.38,-6.37,-6.27,-6.91,-8.22,-7.35,-8.39,-7.87,-9.72,-8.65)
set.seed(123)
M1 <- exp(outer(k1,b)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
fit <- CAES(x=x,M1=M1,M2=M2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CBDCS *CBD with cohort model*

Description

Fits and forecasts mortality rates using CBD with cohort model.

Usage

```
CBDCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).

jumpoff — if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with cohort (M6) model is specified as

$ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \gamma_{t-x} + \epsilon_{x,t}.$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, and γ_c. Constraints include sum of γ_c is zero and sum of $c\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class CBDCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. North American Actuarial Journal, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
fit <- CBDCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CBDQCS — *CBD with curvature and cohort model*

Description

Fits and forecasts mortality rates using CBD with curvature and cohort model.

Usage

```
CBDQCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with curvature and cohort (M7) model is specified as

$ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \kappa_{3,t}((x - \bar{x})^2 - \sigma^2) + \gamma_{t-x} + \epsilon_{x,t}.$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, $\kappa_{3,t}$, and γ_c. Constraints include sum of γ_c is zero, sum of $c\gamma_c$ is zero, and sum of $c^2\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class CBDQCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. North American Actuarial Journal, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
fit <- CBDQCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CBDS *CBD model*

Description

Fits and forecasts mortality rates using CBD model.

Usage

```
CBDS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD (M5) model is specified as

$ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \epsilon_{x,t}.$

The model is estimated by regression and is forecasted by ARIMA applied to $\kappa_{1,t}$ and $\kappa_{2,t}$. It is designed for ages 50-90.

Value

An object of class CBDS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Cairns, A.J.G., Blake, D., and Dowd, K. (2006). A two-factor model for stochastic mortality with parameter uncertainty: Theory and calibration. Journal of Risk and Insurance, 73(4), 687-718.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
fit <- CBDS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CC *Mortality curve calculation*

Description

Calculates mortality rates given mortality curve parameter values.

Usage

```
CC(
  x,
  par,
 curve = c("gompertz", "makeham", "oppermann", "thiele", "wittsteinbumsted", "perks",
   "weibull", "vandermaen", "beard", "heligmanpollard", "rogersplanck", "siler",
    "martinelle", "thatcher")
)
```

Arguments

x	vector of ages.
par	vector of mortality curve parameter values.
curve	name of mortality curve (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher).

Details

See `MC()` for more details of different mortality curves.

Value

Calculated mortality rates based on selected mortality curve and parameter values.

Examples

```
CC(x=60:89,par=c(0.0000082,0.10771),curve="gompertz")
```

CFM2S *Common factor model with global age pattern*

Description

Fits and forecasts mortality rates of two populations using common factor model with global age pattern.

Usage

```
CFM2S(x, M1, M2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M1	matrix of mortality rates of population 1 (rows as years and columns as ages).
M2	matrix of mortality rates of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model with global age pattern is specified as

$ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_x \kappa_{t,i} + \epsilon_{x,t,i}$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of β_x is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class CFM2S with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, J., Wang, M., Liu, J., and Leung, J.W.Y. (2026). Financial valuation of retirement village via stochastic modelling of disability prevalence rates. ASTIN Bulletin, 56(2), 447-473.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b <- c(-0.00010,0.01195,0.03030,0.02170,0.03690,0.02365,0.02280,0.03850,0.05845,0.04415,
0.04185,0.05175,0.03670,0.04195,0.04090,0.02775,0.04990,0.02865,0.03935,0.03820,
0.04000,0.02790,0.03705,0.03370,0.02940,0.02850,0.03400,0.02310,0.02675,0.03430)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
fit <- CFM2S(x=x,M1=M1,M2=M2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CFMS *Common factor model*

Description

Fits and forecasts mortality rates of two populations using common factor model.

Usage

```
CFMS(x, M1, M2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M1	matrix of mortality rates of population 1 (rows as years and columns as ages).
M2	matrix of mortality rates of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model (CFM) is specified as

$ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_{x,i}\kappa_{t,i} + \epsilon_{x,t,i}$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of $\beta_{x,i}$ is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class CFMS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, N. and Lee, R. (2005). Coherent mortality forecasts for a group of populations: An extension of the Lee-Carter method. Demography, 42(3), 575-594.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
```

```
2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b1 <- c(0.0012,-0.0033,0.0523,0.0161,0.0529,0.0220,0.0312,0.0437,0.0709,0.0444,
0.0398,0.0361,0.0403,0.0396,0.0506,0.0315,0.0428,0.0261,0.0384,0.0388,
0.0300,0.0269,0.0275,0.0256,0.0239,0.0421,0.0314,0.0284,0.0174,0.0314)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
b2 <- c(-0.0014,0.0272,0.0083,0.0273,0.0209,0.0253,0.0144,0.0333,0.0460,0.0439,
0.0439,0.0674,0.0331,0.0443,0.0312,0.0240,0.0570,0.0312,0.0403,0.0376,
0.0500,0.0289,0.0466,0.0418,0.0349,0.0149,0.0366,0.0178,0.0361,0.0372)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b1)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b2)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
fit <- CFMS(x=x,M1=M1,M2=M2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

---

curveS3 *Common S3 methods for mortality curves*

Description

Common S3 methods for objects returned by `MC()`:

coef(object) gives estimated parameter values.

fitted(object) gives fitted values.

predict(object,x) gives estimated mortality rate(s) for given age(s) x.

plot(object) plots data and fitted mortality curve.

deviance(object) gives weighted sum of squared residuals on $ln(m_x)$.

residuals(object) gives residuals on $ln(m_x)$.

ENI *Mortality ensemble interval*

Description

Generates ensemble interval forecast of mortality rates.

Usage

```
ENI(..., width = 0.95, method = 1, wm, nsim = 10, seed = 123)
```

Arguments

...	fitted model objects returned by LCS(), RHS(), APCS(), CBDS(), CBDCS(), CBDQCS(), and / or STARS().
width	coverage probability of interval (default = 0.95).
method	if 1, simple averaging; if 2, weighted averaging; if 3, envelope; if 4, interior trimming (default = 1).
wm	vector of weights for LC, RH, APC, M5, M6, M7, and / or STAR models if method = 2 (default = equal weights).
nsim	number of simulations (default = 10).
seed	seed for random number generator (default = 123).

Details

Ensemble interval forecast is constructed by combining interval forecasts from individual stochastic mortality models using different methods including simple averaging, weighted averaging, envelope, and interior trimming. See LCS(), RHS(), APCS(), CBDS(), CBDCS(), CBDQCS(), and STARS() for more details of different stochastic mortality models.

Value

An object of class ENI with associated S3 methods forecast and plot.

References

Li, J., Wang, M., Liu, J., and Tickle, L. (2025). Ensemble interval forecasts of mortality. Scandinavian Actuarial Journal, 2025(6), 598-616.

Examples

```
x <- 60:69
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962)
b <- c(0.0801,0.0909,0.0948,0.0951,0.0965,0.1014,0.1042,0.1141,0.1110,0.1118)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.60,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.90,-4.03,-4.12,
-5.18,-5.64,-6.00,-6.51,-6.91,-6.90,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=10,byrow=TRUE)+rnorm(300,0,0.035))
fit1 <- LCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit2 <- RHS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit3 <- APCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit4 <- CBDS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit <- ENI(fit1,fit2,fit3,fit4)
forecast::forecast(fit)
plot(fit)
```

ENS *Mortality ensemble*

Description

Fits and forecasts mortality rates using mortality ensemble.

Usage

```
ENS(x, M, wm = rep(1/7, 7), curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
wm	vector of weights for LC, RH, APC, M5, M6, M7, and STAR models (default = 1/7).
curve	name of mortality curve for smoothing ensemble mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

Ensemble forecast is obtained as a weighted average of forecasts from individual stochastic mortality models. See `LCS()`, `RHS()`, `APCS()`, `CBDS()`, `CBDCS()`, `CBDQCS()`, and `STARS()` for more details of different stochastic mortality models.

Value

An object of class ENS with associated S3 methods forecast and plot.

References

Li, J. (2023). A model stacking approach for forecasting mortality. North American Actuarial Journal, 27(3), 530-545.

Examples

```
x <- 60:69
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962)
b <- c(0.0801,0.0909,0.0948,0.0951,0.0965,0.1014,0.1042,0.1141,0.1110,0.1118)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.60,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.90,-4.03,-4.12,
-5.18,-5.64,-6.00,-6.51,-6.91,-6.90,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=10,byrow=TRUE)+rnorm(300,0,0.035))
fit <- ENS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
forecast::forecast(fit)
plot(fit)
```

FCS *Mortality model fitting*

Description

Fits and forecasts mortality rates using different stochastic mortality models with different estimation methods.

Usage

```
FCS(
  x,
  M,
  model = c("LC", "RH", "APC", "CBD", "CBDC", "CBDQC", "STAR"),
  curve = NULL,
  h = 10,
  jumpoff = 1
)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
model	name of stochastic mortality model (including LC, RH, APC, CBD, CBDC, CBDQC, and STAR).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).

h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

See LCS(), RHS(), APCS(), CBDS(), CBDCS(), CBDQCS(), and STARS() for more details of different stochastic mortality models.

Value

An object of class based on selected stochastic mortality model with associated S3 methods coef, forecast, plot, and residuals.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- FCS(x=x,M=M,model="LC",curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

LCLS *Lee-Carter model for limited data*

Description

Fits and forecasts mortality rates using Lee-Carter model with sparse data in irregular years.

Usage

```
LCLS(x, t, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
t	vector of years.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Lee-Carter (LC) model is specified as

$ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \epsilon_{x,t}$.

The model is estimated by singular value decomposition and is forecasted by random walk with drift applied to κ_t. Constraints include sum of β_x is one and sum of κ_t is zero. It can be applied to whole age range.

Value

An object of class LCLS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Li, N., Lee, R., and Tuljapurkar, S. (2004). Using the Lee-Carter method to forecast mortality for populations with limited data. International Statistical Review, 72(1), 19-36.

Examples

```
x <- 60:89
t <- c(1991,1996,2001,2006,2011:2020)
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,8.21,
3.27,-1.03,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
```

```
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=14,ncol=30,byrow=TRUE)+rnorm(420,0,0.035))
fit <- LCLS(x=x,t=t,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

LCS *Lee-Carter model*

Description

Fits and forecasts mortality rates using Lee-Carter model.

Usage

```
LCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Lee-Carter (LC) model is specified as

$ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \epsilon_{x,t}$.

The model is estimated by singular value decomposition and is forecasted by ARIMA applied to κ_t. Constraints include sum of β_x is one and sum of κ_t is zero. It can be applied to whole age range.

Value

An object of class LCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Lee, R.D. and Carter, L.R. (1992). Modeling and forecasting U.S. mortality. Journal of the American Statistical Association, 87(419), 659-671.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- LCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

MC *Mortality curve fitting*

Description

Fits parametric mortality curves to observed mortality rates using multiple optimisation strategies.

Usage

```
MC(x, m, curve = .curves, w = rep(1, length(x)))
```

Arguments

x	vector of ages.
m	vector of mortality rates.
curve	name of mortality curve (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive).
w	vector of weights (default = 1).

Details

User can choose one of the following 14 mortality curves (with their suitable age ranges in brackets):

- Gompertz (1825) (ages 30-90) - $m_x = Be^{Cx}$
- Makeham (1860) (ages 20-90) - $m_x = A + Be^{Cx}$
- Oppermann (1870) (ages 0-20) - $m_x = \frac{A}{\sqrt{x+1}} + B + C\sqrt{x+1}$
- Thiele (1871) (ages 0-90) - $m_x = A_1e^{-B_1x} + A_2e^{-0.5B_2(x-C)^2} + A_3e^{B_3x}$
- Wittstein & Bumsted (1883) (ages 0-90) - $m_x = \frac{A^{-(Bx)^N}}{B} + A^{-(M-x)^N}$
- Perks (1932) (ages 20-100+) - $m_x = \frac{A+BC^x}{1+DC^x}$
- Weibull (1939) (ages 50-100+) - $m_x = Bx^C$
- Van der Maen (1943) (ages 80-100+) - $m_x = A + Bx + Cx^2 + \frac{I}{N-x}$
- Beard (1971) (ages 50-100+) - $m_x = \frac{Ae^{Cx}}{1+Be^{Cx}}$
- Heligman & Pollard (1980) (ages 0-100+) - $m_x = A^{(x+B)^C} + De^{-E(ln(x)-ln(F))^2} + \frac{GH^x}{1+GH^x}$
- Rogers & Planck (1983) (ages 0-100+) - $m_x = A_0 + A_1e^{-Ax} + A_2e^{-B(x-U)-e^{-C(x-U)}} + A_3e^{Dx}$
- Siler (1983) (ages 0-90) - $m_x = A_1e^{-B_1x} + A_2 + A_3e^{B_3x}$
- Martinelle (1987) (ages 20-100+) - $m_x = \frac{A+Be^{Cx}}{1+De^{Cx}} + Ee^{Cx}$
- Thatcher (1999) (ages 20-100+) - $m_x = A + \frac{Be^{Cx}}{1+De^{Cx}}$

The mortality curves are fitted by employing multiple optimisation strategies simultaneously (including PORT routines, Nelder-Mead method, and Levenberg-Marquardt algorithm) and selecting one with smallest weighted least squares on $ln(m_x)$. Constrained parameterisations offer traditional interpretability while unconstrained parameterisations provide increased flexibility.

Value

An object of class based on selected mortality curve with associated S3 methods coef, fitted, predict, plot, deviance, and residuals.

References

Gompertz, B. (1825). On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. Philosophical Transactions of the Royal Society of London, 115(1825), 513-583.

Makeham, W.M. (1860). On the law of mortality and the construction of annuity tables. Journal of the Institute of Actuaries, 8(6), 301-310.

Oppermann, L.H.F. (1870). On the graduation of life tables, with special application to the rate of mortality in infancy and childhood. The Institute of Actuaries.

Thiele, T.N. (1871). On a mathematical formula to express the rate of mortality throughout the whole of life, tested by a series of observations made use of by the Danish Life Insurance Company of 1871. Journal of the Institute of Actuaries and Assurance Magazine, 16(5), 313-329.

Wittstein, T. and Bumsted, D.A. (1883). The mathematical law of mortality. Journal of the Institute of Actuaries and Assurance Magazine, 24(3), 153-173.

Perks, W. (1932). On some experiments in the graduation of mortality statistics. Journal of the Institute of Actuaries, 63(1), 12-57.

Weibull, W. (1951). A statistical distribution function of wide applicability. Journal of Applied Mechanics, 18(3), 293-297.

Beard, R.E. (1971). Some aspects of theories of mortality, cause of death analysis, forecasting and stochastic processes. Biological Aspects of Demography, 57-68.

Heligman, L. and Pollard, J.H. (1980). The age pattern of mortality. Journal of the Institute of Actuaries, 107(1), 49-80.

Rogers, A. and Planck, F. (1983). Model: A general program for estimating parametrized model schedules of fertility, mortality, migration, and marital and labor force status transitions. IIASA Working Paper WP-83-102.

Siler, W. (1983). Parameters of mortality in human populations with widely varying life spans. Statistics in Medicine, 2(3), 373-380.

Martinelle, S. (1987). A generalized Perks formula for old-age mortality. Statistiska Centralbyran.

Thatcher, A.R. (1999). The long-term pattern of adult mortality and the highest attained age. Journal of the Royal Statistical Society Series A, 162(1), 5-43.

Tabeau, E. (2001). A review of demographic forecasting models for mortality. Forecasting Mortality in Developed Countries. European Studies of Population, Vol 9.

Examples

```
x <- 60:89
set.seed(123); m <- 0.0000082*exp(0.10771*c(60:89)+rnorm(30,0,0.1))
fit <- MC(x=x,m=m,curve="gompertz")
coef(fit)
fitted(fit)
predict(fit,62.5)
plot(fit)
deviance(fit)
residuals(fit)
```

`modelS3` *Common S3 methods for stochastic mortality models*

Description

Common S3 methods for objects returned by `LCS()`, `RHS()`, `APCS()`, `CBDS()`, `CBDCS()`, `CBDQCS()`, `STARS()`, `ENS()`, `ENI()`, `CFMS()`, `CFM2S()`, `CAES()`, `PLCS()`, `PRHS()`, `PAPCS()`, `PCBDS()`, `PCBDCS()`, `PCBDQCS()`, `PCFMS()`, `PCFM2S()`, `PCAES()`, and `LCLS()`.

coef(object) gives estimated parameter values.

forecast::forecast(object) gives mortality forecasts (which = 1 for smoothed (default); which = 2 for raw) (or ensemble interval forecasts for `ENI()`).

plot(object) plots estimated parameter values, standardised residuals heatmap, data, and mortality forecasts smoothed by selected mortality curve (or ensemble interval forecasts for `ENI()`).

residuals(object) gives standardised residuals on $ln(m_{x,t})$ or $D_{x,t}$.

simulate(object,nsim,seed) gives simulated future mortality rates (raw).

`mortalitymatrix` *Convert mortality data to matrix*

Description

Converts mortality rates stored as a matrix, data frame, or demogdata object into required matrix format.

Usage

```
mortalitymatrix(data, series = 1)
```

Arguments

`data`	mortality data.
`series`	mortality series index for demogdata objects.

Value

A numeric matrix of mortality data.

Examples

```
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
Data <- as.data.frame(M)
mortalitymatrix(Data)
```

PAPCS *Poisson age-period-cohort model*

Description

Fits and forecasts mortality rates using age-period-cohort model with Poisson assumption.

Usage

```
PAPCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The age-period-cohort (APC) model with Poisson assumption is specified as

$ln(m_{x,t}) = \alpha_x + \kappa_t + \gamma_{t-x}$ and $D_{x,t}\ Poisson(E_{x,t}m_{x,t})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c. Constraints include sum of κ_t is zero and and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class PAPCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Haberman, S. and Renshaw, A. (2011). A comparative study of parametric mortality projection models. Insurance: Mathematics and Economics, 48(1), 35-55.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- rep(1/30,30)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PAPCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCAES *Poisson common age effect model*

Description

Fits and forecasts mortality rates of two populations using common age effect model with Poisson assumption.

Usage

```
PCAES(x, D1, D2, E1, E2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D1	matrix of death counts of population 1 (rows as years and columns as ages).
D2	matrix of death counts of population 2 (rows as years and columns as ages).
E1	matrix of mid-year exposures of population 1 (rows as years and columns as ages).
E2	matrix of mid-year exposures of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2,

	vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common age effect (CAE) model with Poisson assumption is specified as

$ln(m_{x,t,i}) = \alpha_{x,i} + \beta_x \kappa_{t,i}$ and $D_{x,t,i}\ Poisson(E_{x,t,i} m_{x,t,i})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{t,i}$. Constraints include sum of β_x is one and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class PCAES with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Kleinow, T. (2015). A common age effect model for the mortality of multiple populations. Insurance: Mathematics and Economics, 63(C), 147-152.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
b <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
k1 <- c(8.68,8.34,7.99,6.87,8.18,5.73,4.83,5.20,2.74,3.22,
2.99,1.59,1.67,-0.65,-0.39,-1.07,-0.95,-2.78,-3.46,-2.45,
-4.12,-4.66,-4.98,-4.58,-6.30,-4.39,-5.56,-6.52,-8.26,-6.92)
k2 <- c(11.81,11.01,10.59,10.40,9.75,8.15,6.07,6.45,4.60,4.57,
4.15,1.49,1.77,-1.08,-1.44,-0.96,-1.66,-2.25,-4.67,-4.62,
-4.38,-6.37,-6.27,-6.91,-8.22,-7.35,-8.39,-7.87,-9.72,-8.65)
set.seed(123)
M1 <- exp(outer(k1,b)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
E1 <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
```

```
E2 <- E1
D1 <- round(E1*M1)
D2 <- round(E2*M2)
fit <- PCAES(x=x,D1=D1,D2=D2,E1=E1,E2=E2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCBDCS *Poisson CBD with cohort model*

Description

Fits and forecasts mortality rates using CBD with cohort model with Poisson assumption.

Usage

```
PCBDCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with cohort (M6) model with Poisson assumption is specified as

$ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \gamma_{t-x}$ and $D_{x,t}\ Poisson(E_{x,t}m_{x,t})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, and γ_c. Constraints include sum of γ_c is zero and sum of $c\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class PCBDCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. North American Actuarial Journal, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PCBDCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCBDQCS *Poisson CBD with curvature and cohort model*

Description

Fits and forecasts mortality rates using CBD with curvature and cohort model with Poisson assumption.

Usage

```
PCBDQCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with curvature and cohort (M7) model with Poisson assumption is specified as

$ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \kappa_{3,t}((x - \bar{x})^2 - \sigma^2) + \gamma_{t-x}$ and $D_{x,t}$ $Poisson(E_{x,t}m_{x,t})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, $\kappa_{3,t}$, and γ_c. Constraints include sum of γ_c is zero, sum of $c\gamma_c$ is zero, and sum of $c^2\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class PCBDQCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. North American Actuarial Journal, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PCBDQCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCBDS *Poisson CBD model*

Description

Fits and forecasts mortality rates using CBD model with Poisson assumption.

Usage

```
PCBDS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD (M5) model with Poisson assumption is specified as

$ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x})$ and $D_{x,t}\ Poisson(E_{x,t}m_{x,t})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$ and $\kappa_{2,t}$. It is designed for ages 50-90.

Value

An object of class PCBDS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Cairns, A.J.G., Blake, D., and Dowd, K. (2006). A two-factor model for stochastic mortality with parameter uncertainty: Theory and calibration. Journal of Risk and Insurance, 73(4), 687-718.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PCBDS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCFM2S *Poisson common factor model with global age pattern*

Description

Fits and forecasts mortality rates of two populations using common factor model with global age pattern with Poisson assumption.

Usage

```
PCFM2S(x, D1, D2, E1, E2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D1	matrix of death counts of population 1 (rows as years and columns as ages).
D2	matrix of death counts of population 2 (rows as years and columns as ages).
E1	matrix of mid-year exposures of population 1 (rows as years and columns as ages).
E2	matrix of mid-year exposures of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model with global age pattern with Poisson assumption is specified as $ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_x \kappa_{t,i}$ and $D_{x,t,i}\ Poisson(E_{x,t,i} m_{x,t,i})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of β_x is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class PCFM2S with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, J., Wang, M., Liu, J., and Leung, J.W.Y. (2026). Financial valuation of retirement village via stochastic modelling of disability prevalence rates. ASTIN Bulletin, 56(2), 447-473.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b <- c(-0.00010,0.01195,0.03030,0.02170,0.03690,0.02365,0.02280,0.03850,0.05845,0.04415,
0.04185,0.05175,0.03670,0.04195,0.04090,0.02775,0.04990,0.02865,0.03935,0.03820,
0.04000,0.02790,0.03705,0.03370,0.02940,0.02850,0.03400,0.02310,0.02675,0.03430)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
E1 <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
E2 <- E1
D1 <- round(E1*M1)
D2 <- round(E2*M2)
fit <- PCFM2S(x=x,D1=D1,D2=D2,E1=E1,E2=E2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCFMS *Poisson common factor model*

Description

Fits and forecasts mortality rates of two populations using common factor model with Poisson assumption.

Usage

```
PCFMS(x, D1, D2, E1, E2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D1	matrix of death counts of population 1 (rows as years and columns as ages).
D2	matrix of death counts of population 2 (rows as years and columns as ages).
E1	matrix of mid-year exposures of population 1 (rows as years and columns as ages).
E2	matrix of mid-year exposures of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model (CFM) with Poisson assumption is specified as

$ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_{x,i}\kappa_{t,i}$ and $D_{x,t,i}\ Poisson(E_{x,t,i}m_{x,t,i})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of $\beta_{x,i}$ is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class PCFMS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, J. (2013). A Poisson common factor model for projecting mortality and life expectancy jointly for females and males. Population Studies, 67(1), 111-126.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b1 <- c(0.0012,-0.0033,0.0523,0.0161,0.0529,0.0220,0.0312,0.0437,0.0709,0.0444,
0.0398,0.0361,0.0403,0.0396,0.0506,0.0315,0.0428,0.0261,0.0384,0.0388,
0.0300,0.0269,0.0275,0.0256,0.0239,0.0421,0.0314,0.0284,0.0174,0.0314)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
b2 <- c(-0.0014,0.0272,0.0083,0.0273,0.0209,0.0253,0.0144,0.0333,0.0460,0.0439,
0.0439,0.0674,0.0331,0.0443,0.0312,0.0240,0.0570,0.0312,0.0403,0.0376,
0.0500,0.0289,0.0466,0.0418,0.0349,0.0149,0.0366,0.0178,0.0361,0.0372)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b1)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b2)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
E1 <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
E2 <- E1
D1 <- round(E1*M1)
D2 <- round(E2*M2)
fit <- PCFMS(x=x,D1=D1,D2=D2,E1=E1,E2=E2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PFCS *Poisson mortality model fitting*

Description

Fits and forecasts mortality rates using different stochastic mortality models with Poisson assumption.

Usage

```
PFCS(
  x,
  D,
  E,
  model = c("LC", "RH", "APC", "CBD", "CBDC", "CBDQC"),
  curve = NULL,
  h = 10,
  jumpoff = 1
)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
model	name of stochastic mortality model (including LC, RH, APC, CBD, CBDC, and CBDQC).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

See PLCS(), PRHS(), PAPCS(), PCBDS(), PCBDCS(), and PCBDQCS() for more details of different stochastic mortality models with Poisson assumption.

Value

An object of class based on selected stochastic mortality model with associated S3 methods coef, forecast, plot, and residuals.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
```

```
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PFCS(x=x,D=D,E=E,model="LC",curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PLCS *Poisson Lee-Carter model*

Description

Fits and forecasts mortality rates using Lee-Carter model with Poisson assumption.

Usage

```
PLCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Lee-Carter (LC) model with Poisson assumption is specified as

$ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t$ and $D_{x,t}$ $Poisson(E_{x,t} m_{x,t})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t. Constraints include sum of β_x is one and sum of κ_t is zero. It can be applied to whole age range.

Value

An object of class PLCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Renshaw, A.E. and Haberman, S. (2006). A cohort-based extension to the Lee–Carter model for mortality reduction factors. Insurance: Mathematics and Economics, 38(3), 556-570.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PLCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PRHS *Poisson Renshaw-Haberman model*

Description

Fits and forecasts mortality rates using Renshaw-Haberman model with Poisson assumption.

Usage

```
PRHS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Renshaw-Haberman (RH) model with Poisson assumption is specified as

$ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \gamma_{t-x}$ and $D_{x,t}\ Poisson(E_{x,t} m_{x,t})$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c. Constraints include sum of β_x is one, sum of κ_t is zero, and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class PRHS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Haberman, S. and Renshaw, A. (2011). A comparative study of parametric mortality projection models. Insurance: Mathematics and Economics, 48(1), 35-55.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
```

```
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PRHS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

RHS *Renshaw-Haberman model*

Description

Fits and forecasts mortality rates using Renshaw-Haberman model.

Usage

```
RHS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Renshaw-Haberman (RH) model is specified as

$ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \gamma_{t-x} + \epsilon_{x,t}$.

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c. Constraints include sum of β_x is one, sum of κ_t is zero, and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class RHS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Haberman, S. and Renshaw, A. (2011). A comparative study of parametric mortality projection models. Insurance: Mathematics and Economics, 48(1), 35-55.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- RHS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

SEN *Lee-Carter model sensitivities*

Description

Calculates sensitivities of Lee-Carter model parameters, mortality forecasts, and derived actuarial quantities to small changes in data.

Usage

```
SEN(
  M,
  x,
  t,
  x1 = NULL,
  t1 = NULL,
  n1 = NULL,
  r = NULL,
  f = c("s", "u", "v", "a", "b", "k", "kf", "mf", "axn", "A1xn"),
  jumpoff = 1,
  mq = 1
)
```

Arguments

M	matrix of central death rates (rows as years, columns as ages).
x	index age of data point (column number of M) being perturbed.
t	index year of data point (row number of M) being perturbed.
x1	index age of output variable (or initial age for annuity or assurance) whose sensitivity is calculated.
t1	index year of output variable whose sensitivity is calculated.
n1	term of annuity or assurance whose sensitivity is calculated.
r	annual interest rate for annuity or assurance whose sensitivity is calculated.
f	type of output variable whose sensitivity is calculated (including s, u, v, a, b, k, kf, mf, axn, A1xn).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed death rates in final year (default = 1).
mq	if 1, sensitivity is calculated with respect to central death rate; if 2, sensitivity is calculated with respect to probability of death (default = 1).

Details

User can choose one of the 10 output variables, including the first singular value (s), first left singular vector (u), first right singular vector (v), alpha (a), beta (b), kappa (k), forecasted kappa (kf), forecasted death rates (mf), expected present value of a term annuity ($1 payable at the end of each year of survival) (axn), and expected present value of a term assurance ($1 payable at the end of the year of death) (A1xn), under the Lee-Carter model, forecasted by applying the random walk with drift to κ_t. Their sensitivities are analytically derived using first-order matrix perturbation theory.

Value

An object of class SEN with associated S3 methods coef and forecast.

References

Li, J. and Haberman, S. (2026). The Lee-Carter model revisited: Perturbation analysis and estimation equivalence. Working Paper.

Examples

```
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
SEN(M,x=10,t=15,f="s")$sensitivity
SEN(M,x=10,t=15,x1=10,f="u")$sensitivity
SEN(M,x=10,t=15,t1=15,f="v")$sensitivity
SEN(M,x=10,t=15,x1=10,f="a")$sensitivity
SEN(M,x=10,t=15,x1=10,f="b")$sensitivity
SEN(M,x=10,t=15,t1=15,f="k")$sensitivity
SEN(M,x=10,t=15,t1=35,f="kf")$sensitivity
SEN(M,x=10,t=15,x1=10,t1=35,f="mf")$sensitivity
SEN(M,x=10,t=15,x1=10,n1=10,r=0.02,f="axn")$sensitivity
SEN(M,x=10,t=15,x1=10,n1=10,r=0.02,f="A1xn")$sensitivity
```

STARS *Spatial-temporal autoregressive model*

Description

Fits and forecasts mortality rates using spatial-temporal autoregressive model.

Usage

```
STARS(x, M, curve = NULL, h = 10)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2,

vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).

h forecast horizon (default = 10).

Details

The spatial-temporal autoregressive (STAR) model is specified as

$ln(M_t) = \mu + Rln(M_{t-1}) + E_t,$

where μ contains intercept parameters and R is banded matrix with non-zero entries in main diagonal and two subdiagonals below that.

The model is estimated by constrained regression and is forecasted via its autoregressive nature. Constraints in R include: all non-zero entries are between zero and one and sum of each row is one. It can be applied to whole age range.

Value

An object of class STARS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Li, H. and Lu, Y. (2017). Coherent forecasting of mortality rates: A sparse vector-autoregression approach. ASTIN Bulletin, 47(2), 563-600.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- STARS(x=x,M=M,curve="makeham",h=30)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

Index

APCS, 2

CAES, 4
CBDCS, 5
CBDQCS, 6
CBDS, 8
CC, 9
CFM2S, 10
CFMS, 11
curveS3, 13

ENI, 13
ENS, 15

FCS, 16

LCLS, 17
LCS, 19

MC, 20
modelS3, 22
mortalitymatrix, 23

PAPCS, 24
PCAES, 25
PCBDCS, 27
PCBDQCS, 28
PCBDS, 29
PCFM2S, 31
PCFMS, 32
PFCS, 34
PLCS, 36
PRHS, 37

RHS, 39

SEN, 40
STARS, 42